

Package ‘OptionPricing’

January 20, 2025

Type Package

Title Option Pricing with Efficient Simulation Algorithms

Version 0.1.2

Maintainer Wolfgang Hormann <hormanngw@yahoo.com>

Description Efficient Monte Carlo Algorithms for the price and the sensitivities of Asian and European Options under Geometric Brownian Motion.

License GPL-2 | GPL-3

Copyright Wolfgang Hormann

NeedsCompilation no

Repository CRAN

Date/Publication 2023-09-16 08:40:08 UTC

Author Wolfgang Hormann [aut, cre],
Kemal Dingec [aut]

Contents

OptionPricing-package	1
AsianCall	2
AsianCall_AppLord	5
BS_EC	6
Index	7

OptionPricing-package *Option Pricing and Greeks Estimation for Asian and European Options*

Description

The Price, Delta and Gamma of European and Asian Options under Geometric Brownian Motion are calculated using the Black-Scholes formula and Efficient Monte Carlo and Randomized Quasi Monte Carlo Algorithms.

Details

The OptionPricing package calculates the Price, Delta and Gamma for European options using the Black-Scholes formula (see [BS_EC](#)). The price, Delta and Gamma for Asian call options under geometric Brownian motion are calculated using a very efficient Monte Carlo and randomized quasi-Monte Carlo algorithm (see [AsianCall](#)). The function [AsianCall_AppLord](#) implements a high-quality approximation for the price of an Asian option.

Author(s)

Kemal Dingec, Wolfgang Hormann

Examples

```
# standard settings for an efficient simulation using QMC and variance reduction
AsianCall(T=1,d=12,K=100,r=0.05,sigma=0.2,S0=100,method=c("best"),
  sampling=c("QMC"),metpar=list(maxiter=100,tol=1.e-14,cvmethod="splitting"),
  sampar=list(nout=50,n=2039,a=1487,baker=TRUE,genmethod="pca"))

# Calculation of the Price of an Asian option using a good approximation
AsianCall_AppLord(T = 1, d = 12, K = 100, r = 0.05, sigma = 0.2, S0 = 100)

# standard settings for an efficient simulation using MC and variance reduction
AsianCall(T=1,d=12,K=170,r=0.05,sigma=0.2,S0=100,method="best",
  sampling="MC",metpar=list(maxiter=100,tol=1.e-14,np=1000),
  sampar=list(n=10^5))
# Calculation of the approximate price, a bit different to the above result
AsianCall_AppLord(T = 1, d = 12, K = 170, r = 0.05, sigma = 0.2, S0 = 100)

# Calculation of the Price of an Asian option using a good approximation
AsianCall_AppLord(T = 1, d = 12, K = 100, r = 0.05, sigma = 0.2, S0 = 100)

#Price, Delta and Gamma of European options using Black-Scholes
BS_EC(K=100, r = 0.05, sigma = 0.2, T = 0.25, S0 = 100)
BS_EP(K=100, r = 0.05, sigma = 0.2, T = 0.25, S0 = 100)
```

AsianCall

Calculates the Price, Delta and Gamma of an Asian Option

Description

Prices arithmetic average Asian Call options under geometric Brownian motion. It also estimates the sensitivities Delta and Gamma.

Usage

```
AsianCall(T=1,d=12,K=100,r=0.05,sigma=0.2,S0=100,method=c("best","naive"),
  sampling=c("QMC","MC"),
  metpar=list(maxiter=100,tol=1.e-14,cvmethod="splitting"),
```

```
sampar=list(nout=50,seq.type="korobov",n=2039,a=1487,
            baker=TRUE,genmethod="pca"))
```

Arguments

T	time to maturity (in years)
d	number of control points
K	strike price
r	risk free interest rate
sigma	volatility
S0	starting stockprice
method	selects the simulation method; method "best" uses a variance reduction method based on effective control variates and conditional Monte Carlo and is very effective. method "naive" is mainly provided for comparison purposes.
sampling	sampling QMC uses the Quasi Monte Carlo method Korobov lattice for the simulation. sampling MC uses standard Monte Carlo for the simulation.
metpar	list holding extra parameters related to the simulation method For method="best": maxiter= maximal no of iterations for Newton method tol= error tolerance for Newton method for sampling="QMC": cvmethod=c("splitting","direct") NOT necessary for method = "naive" "splitting" ... estimates CV coefficients using lm with bootstrap "direct" ... estimates CV coefficients using lm and the full sample for sampling="MC": np ... sample size for pilot run for CV; NOT necessary for method = "naive"
sampar	list holding several parameters related to the sampling method; for sampling="MC" the list sampar only contains the total samplesize n; for sampling="QMC" the list sampar contains the elements: nout number of independent "randomized" copies of the Korobov lattice n number of points of the Korobov lattice a important constant for the construction of the Korobov lattice baker TRUE/FALSE, indicates if Baker transform should be used for making the integrand periodic genmethod = c("pca", "std", "pcamain", "lt", "ltpca"), note that for method=="naive" only genmethod=c("pca", "std") can be used. genmethod="pca" principal component analysis genmethod="std" standard genmethod="pcamain" use only first dirnum main directions of the PCA

genmethod="lt" uses a transform for the first dirnum
 genmethod="ltpca" combination of lt with pca
 dirnum number of main directions, only used for genmethod="pcamain" or
 "lt"

Details

Method best (see the reference Dingec and Hormann below) is a very efficient simulation algorithm using multiple Control Variates and conditional MonteCarlo to calculate the the price, delta and gamma of Asian call options under geometric Brownian motion. It is especially effective when QMC is selected as sampling method. As QMC method Korobov Lattice rules are used. For good parameter values see Table 1 of (L'Ecuyer, Lemieux).

Value

returns a matrix holding the price and greeks. The estimated Asian Call price and its estimated delta and gamma form the first column vector, the respective 95 percent error bounds are given in the second column.

Author(s)

Kemal Dingec, Wolfgang Hormann

References

K. D. Dingec and W. Hormann. Improved Monte Carlo and Quasi-Monte Carlo Methods for the Price and the Greeks of Asian Options, Proceedings of the 2014 Winter Simulation Conference A. Tolk, S. D. Diallo, I. O. Ryzhov, L. Yilmaz, S. Buckley, and J. A. Miller, eds.

L'Ecuyer, P., and C. Lemieux. 2000. Variance Reduction via Lattice Rules. Management Science 46 (9): 1214-1235.

See Also

[OptionPricing-package](#)

Examples

```
# standard settings for an efficient simulation using QMC and variance reduction
AsianCall(T=1,d=12,K=100,r=0.05,sigma=0.2,S0=100,method="best",
  sampling="QMC",metpar=list(maxiter=100,tol=1.e-14,cvmethod="splitting"),
  sampar=list(nout=50,n=2039,a=1487,baker=TRUE,genmethod="pca"))
```

```
# efficient Monte Carlo version of the above simulation
AsianCall(T=1,d=12,K=100,r=0.05,sigma=0.2,S0=100,method="best",
  sampling="MC",metpar=list(maxiter=100,tol=1.e-14,np=1000),
  sampar=list(n=10^5))
```

```
# simple QMC version without variance reduction
```

```
AsianCall(T=1,d=12,K=100,r=0.05,sigma=0.2,S0=100,method="naive",
```

```

    sampling="QMC",
    sampar=list(nout=50,n=2039,a=1487,baker=TRUE,genmethod="pca"))

# naive Monte Carlo version
AsianCall(T=1,d=12,K=100,r=0.05,sigma=0.2,S0=100,method="naive",
    sampling="MC",sampar=list(n=10^5))

```

AsianCall_AppLord *Asian Options - Approximation*

Description

The price of an arithmetic average Asian option is computed using the approximation method of Lord.

Usage

```
AsianCall_AppLord(T=1, d=12, K=100, r=0.05, sigma=0.1, S0=100, all=TRUE)
```

Arguments

T	T time to maturity (in years)
d	d number of controll points
K	K strike price
r	r risk free interest rate
sigma	sigma volatility (yearly)
S0	S0 starting stockprice
all	all TRUE means that the full Asian Call option price is approximated

Details

AsianCall_AppLord() uses a sophisticated approximation of Lord (2006).

Value

returns the approximate price.

Author(s)

Kemal Dingec, Wolfgang Hormann

References

Lord, R., Partially Exact and Bounded Approximations for Arithmetic Asian Options, Journal of Computational Finance, Vol. 10, No. 2, pp. 1-52, 2006

See Also

[OptionPricing-package](#)

Examples

```
AsianCall_AppLord(T = 1, d = 12, K = 100, r = 0.05, sigma = 0.25, S0 = 100, all = TRUE)
```

 BS_EC

Black-Scholes Formula for European Call and Put

Description

Calculates the Price, Delta and Gamma of an European Call or Put option using the Black-Scholes formula.

Usage

```
BS_EC( T = 0.25, K = 100, r = 0.05, sigma = 0.2, S0 = 100 )
BS_EP( T = 0.25, K = 100, r = 0.05, sigma = 0.2, S0 = 100 )
```

Arguments

T	time to maturity (in years)
K	Strike Price
r	risk-free interest rate
sigma	yearly volatility
S0	Starting Stock Price

Value

Returns a vector containing the option price, Delta and Gamma

Author(s)

Wolfgang Hormann

See Also

[OptionPricing-package](#)

Examples

```
BS_EC(K=100, r = 0.05, sigma = 0.2, T = 0.25, S0 = 100)
BS_EP(K=100, r = 0.05, sigma = 0.2, T = 0.25, S0 = 100)
```

Index

* package

OptionPricing-package, 1

AsianCall, 2, 2

AsianCall_AppLord, 2, 5

BS_EC, 2, 6

BS_EP (BS_EC), 6

OptionPricing (OptionPricing-package), 1

OptionPricing-package, 1